

Scaling and memory in the return intervals of energy dissipation rate in three-dimensional fully developed turbulence

Chuang Liu,^{1,2} Zhi-Qiang Jiang,^{1,3} Fei Ren,^{1,2,4} and Wei-Xing Zhou^{1,2,3,4,5,*}

¹*School of Business, East China University of Science and Technology, Shanghai 200237, China*

²*Engineering Research Center of Process Systems Engineering (Ministry of Education), East China University of Science and Technology, Shanghai 200237, China*

³*School of Science, East China University of Science and Technology, Shanghai 200237, China*

⁴*Research Center for Econophysics, East China University of Science and Technology, Shanghai 200237, China*

⁵*Research Center on Fictitious Economics & Data Science, Chinese Academy of Sciences, Beijing 100080, China*

(Dated: December 31, 2008)

We study the statistical properties of return intervals r between successive energy dissipation rates above a certain threshold Q in three-dimensional fully developed turbulence. We find that the distribution function $P_Q(r)$ scales with the mean return interval R_Q as $P_Q(r) = R_Q^{-1} f(r/R_Q)$ except for $r = 1$, where the scaling function $f(x)$ has two power-law regimes. The return intervals are short-term and long-term correlated and possess multifractal nature. The Hurst index of the return intervals decays exponentially against R_Q , predicting that rare extreme events with $R_Q \rightarrow \infty$ are also long-term correlated with the Hurst index $H_\infty = 0.639$.

PACS numbers: 47.27.Jv, 47.53.+n, 05.45.Tp, 89.75.Da

Extreme events are ubiquitous in nature and society and understanding their dynamics is of crucial importance [1, 2]. However, extreme events are usually rare, which makes it difficult to investigate their occurrence properties. Recently, there is an increasing interest in the study of return intervals or reoccurrence times r between successive events above (or below) some threshold Q , aiming at unveiling the laws governing the occurrence of extreme events by studying the statistics of the return intervals for increasing threshold Q .

Recent studies show that the long-term correlation structure has essential influence on the statistics of return intervals [3]. For long-term power-law correlated monofractal records with exponent γ , numerical analysis illustrates that the distribution density function of the return intervals follows a stretched exponential $P_Q(r) \sim \exp[-b(r/R_Q)^\gamma]$ with the same exponent γ and the return intervals are also long-term correlated, again with the same exponent γ , where R_Q is the mean return interval associated with threshold Q [4, 5, 6]. Uncorrelated records with $\gamma = 1$ is a special case, whose return intervals are exponentially distributed. When $0 < \gamma \leq 1$, theoretical analysis with certain approximation shows that the distribution of return interval is Weibull: $P_Q(r) \sim r^{\gamma-1} \exp(-cr^\gamma)$ [7, 8]. This seems consistent with recent numerical results [9].

For multifractal records in the presence or absence of linear correlations, extensive simulations based on the multiplicative random cascade (MRC) model [10] and the multifractal random walk (MRW) model [11] unveil that the return intervals have a power-law decay in the distribution and are long-term correlated governed by power laws whose exponents depend explicitly on the threshold Q , and the conditional return intervals increase

as a power-law function of the previous return interval [12, 13, 14]. These results are of particular interest since a variety of time series exhibit multifractal nature. For instance, the returns of common stocks can be well modeled by the multifractal random walk [15, 16], and the statistics of the associated return intervals are found to comply with the numerical prediction [12, 14], which can be used to significantly improve risk estimation [12].

Another important issue of multifractal records concerns the possible scaling behavior of the return interval distributions $P_Q(r)$ over different thresholds Q . Numerical simulations of MRC and MRW time series find no evidence of such scaling [12, 13, 14]. However, empirical return interval analysis of financial volatility gives miscellaneous results. Several studies reported that there is a scaling law in the return interval distributions [17, 18, 19, 20, 21], while others argued that the cumulative distributions of return intervals had systematic deviations from scaling and showed multiscaling behaviors [22, 23, 24, 25].

In this Letter, we perform return interval analysis of the energy dissipation rate in three-dimensional fully developed turbulence based on a high-Reynolds turbulence data set collected at the S1 ONERA wind tunnel by the Grenoble group from LEGI [26]. The size of the velocity time series $\{v_i : i = 1, 2, \dots, N\}$ is about 1.73×10^7 . Using Taylor's frozen flow hypothesis which replaces a spatial variation of the fluid velocity by a temporal variation measured at a fixed location, the rate of kinetic energy dissipation at position i is $\epsilon_i \sim [(v_{i+1} - v_i)/\delta_\ell]^2$, where δ_ℓ is the resolution (translated in spatial scale) of the measurements. The energy dissipation rate time series exhibit multifractal nature [27] and its Hurst index is $H = 1 - \gamma/2 = 0.81$. Contrary to previous studies,

we find that the return interval distributions show two power-law regimes and collapse onto a single curve for different thresholds Q . The scaling phenomenon is also observed for conditional interval distributions.

We have calculated the return interval time series r_i for different thresholds Q , which can be mapped nonlinearly to the mean return intervals R_Q . Logarithmic binning is adopted to construct the distribution density functions $P_Q(r)$. In order to ensure that the bins cover the whole r -axis, we use the following procedure. First, the interval $[1, \max(r_i)]$ is partitioned logarithmically into $n - 1$ subintervals whose edges are $x_1 < x_2 < \dots < x_n$. Then we obtain the sequence $y_i = [x_i]$, where $[x_i]$ is a round function of x_i . We discard duplicate integers in the y_i sequence and obtain a new sequence w_j . The edge sequence of the bins are determined by $\{e_i\} = \{0.5, \{w_i + 0.5\}\}$. For each bin $(e_i, e_{i+1}]$, the empirical density function can be calculated by

$$P_Q(r_i) = \frac{\#(e_i < r < e_{i+1})}{\#(r > 0)} \frac{1}{e_{i+1} - e_i}, \quad (1)$$

where $r_i = (e_i + e_{i+1})/2$ and $\#()$ is the number of return intervals that satisfies the condition in the parenthesis. The empirical distribution of the return intervals is depicted in Fig. 1 for $R_Q = 50, 150$ and 500 . We find that the three distributions collapse onto a single curve except for $r = 1$, in remarkable contrast to the simulation results for MRC and MRW time series. Two power-law regimes are observed

$$R_Q P_Q(r) \sim \begin{cases} A_1 (r/R_Q)^{-\delta_1} & \text{if } 1 < r < r_c \\ A_2 (r/R_Q)^{-\delta_2} & \text{if } r > r_c \end{cases}, \quad (2)$$

where $A_1 = 0.107$ and $A_2 = 33.4$ are prefactors, the crossover return interval $r_c \approx 7R_Q$, and $\delta_1 = 0.987 \pm 0.013$ and $\delta_2 = 3.88 \pm 0.09$. For the shuffled data, the $R_Q P_Q(r)$ curves collapse to a single exponential curve.

In risk estimation, a quantity of great interest is the probability $W_Q(\Delta t, t)$ that an extreme event occurs after a short time $\Delta t \ll t$ from now on, conditioned that the time elapsed t after the occurrence of the previous extreme event [12]:

$$W_Q(\Delta t|t) = \frac{\int_t^{t+\Delta t} P_Q(t) dt}{\int_t^\infty P_Q(t) dt}. \quad (3)$$

When $t > r_c$, simple algebraic manipulation leads to

$$W_Q(\Delta t|t) \simeq (\delta_2 - 1) \Delta t / t. \quad (4)$$

The probability $W_Q(\Delta t|t)$ is found to be proportional to Δt and inversely proportional to t . An intriguing feature is that $W_Q(\Delta t|t)$ is independent of the threshold Q , which is a direct consequence of the scaling behavior of




FIG. 1: (color online). Log-log plot of the scaled distribution $R_Q P_Q(r)$ as a function of r/R_Q for three different values of R_Q . The straight lines are the best fits to power laws. The power-law exponents are $\delta_1 = 0.987 \pm 0.013$ and $\delta_2 = 3.88 \pm 0.09$. For the shuffled data, the $R_Q P_Q(r)$ curves collapse to a same exponential curve, as expected.

$P_Q(r)$ shown in Fig. 1. When $t < r_c$, we obtain that

$$\begin{aligned} W_Q(\Delta t|t) &\approx \frac{(\delta_1 - 1) \left(\frac{t}{R_Q}\right)^{-\delta_1} \frac{\Delta t}{R_Q}}{\left(\frac{t}{R_Q}\right)^{1-\delta_1} + \frac{A_2}{A_1} \frac{\delta_1 - 1}{\delta_2 - 1} \left(\frac{r_c}{R_Q}\right)^{1-\delta_2} - \left(\frac{r_c}{R_Q}\right)^{1-\delta_1}} \\ &\approx \frac{(\delta_1 - 1) \left(\frac{t}{R_Q}\right)^{-\delta_1} \frac{\Delta t}{R_Q}}{\left(\frac{t}{R_Q}\right)^{1-\delta_1} - 1.0308}. \end{aligned} \quad (5)$$

We find that $W_Q(\Delta t|t)$ is proportional to Δt . However, $W_Q(\Delta t|t)$ also depends on R_Q .

In order to test the memory effects of the return intervals, we first investigate the conditional PDF $P_Q(r|r_0)$, which is the distribution of return intervals immediately after r_0 . To gain better statistics, we study $P_Q(r|r_0)$ for a range of r_0 rather than individual r_0 values. For each threshold Q or R_Q , the return intervals sequence are sorted in an increasing order and then divided into eight groups $G_1, \dots, G_8$ with approximately equal size. An empirical conditional distribution is determined for each r_0 group. Figure 2 shows $P_Q(r|r_0)$ for $r_0 \in G_1$ and $r_0 \in G_8$. For each group, the three distributions collapse onto a single curve, indicating evident scaling behavior. The figure shows that the probability of finding small (large) r in G_1 is enhanced (decreased) compared with G_8 . This discrepancy in the two groups of distributions unveils the memory effect that large (small) return intervals tend to follow large (small) return intervals. This is true since the distributions associated with different G_i should not exhibit significant discrepancy if there is no memory in the return intervals [17].

The memory effect in the conditional distribution $P_Q(r|r_0)$ can also be illustrated by the mean conditional return interval $\langle r|r_0 \rangle$. If there is no memory in the return




FIG. 2: (color online). Scaling and memory in the conditional distributions. The scaled conditional distribution $R_Q P_Q(r|r_0)$ is plotted as a function of r/R_Q with r_0 in G_1 and G_8 for three different values of R_Q .

intervals, $\langle r|r_0 \rangle$ does not depend on r_0 and is equal to R_Q . Figure 3 plots the mean conditional return interval $\langle r|r_0 \rangle$ as a function of r_0 for $R_Q = 50, 150$ and 500 . It is shown that $\langle r|r_0 \rangle$ is a power-law function of r_0 when r_0 is larger than certain value and the power-law exponent decreases with R_Q , which is consistent with many simulational and empirical studies. Also shown in Fig. 3 is the mean conditional return interval of the shuffled energy dissipation rate, which does not depend on r_0 .




FIG. 3: (color online). Scaled mean conditional return interval $\langle r|r_0 \rangle / R_Q$ versus r_0 / R_Q of return interval sequence in turbulence for various R_Q . The filled symbols represent the results for the shuffled records without memory.

We now study the long-term correlation in the return intervals using the multifractal detrended fluctuation analysis (MFDFA), which is able to extract long-term power-law correlation in non-stationary time series [28, 29, 30]. The MFDFA considers the cumulative time series $R_i = \sum_{i=1}^m (r_i - \langle r \rangle)$, which is partitioned into N_s disjoint boxes with the same size s . In each box k , the local trend is removed from the subseries by a poly-

nomial function and the local rms fluctuation $f_k(s)$ is determined. The overall detrended fluctuation is calculated by

$$F_q(s) = \left\{ \frac{1}{N_s} \sum_{k=1}^{N_s} [f_k(s)]^q \right\}^{1/q}. \quad (6)$$

By varying the value of s , one can expect the detrended fluctuation function $F_q(s)$ scales with the size s :

$$F_q(s) \sim s^{h(q)}, \quad (7)$$

where $h(q)$ is the generalized Hurst index. The return interval series possesses multifractal nature if and only if $h(q)$ is a nonlinear function of q . When $q = 2$, $h(2)$ is nothing but the Hurst index H and the MFDFA reduces to the DFA. The Hurst index H is related to the autocorrelation exponent γ by $\gamma = 2 - 2H$.

We first investigate the linear long-term correlation property of the return intervals using DFA. The dependence of the fluctuation function F_2 is drawn in Fig. 4 against s for $R_Q = 50, 150$, and 500 . In all cases, we find nice power-law relation and the scaling range decreases with the increase of R_Q . The inset shows the dependence of the Hurst index with respect to R_Q , which has an exponential decay:

$$H = H_\infty + be^{-R_Q/R_c} = 0.639 + 0.158e^{-R_Q/69.9}, \quad (8)$$

where $R_c = 69.9$ is the characteristic scale. For extreme events with very large Q , R_Q tends to infinity, and the Hurst index can be predicted as $H = H_\infty = 0.639$. This implies that the return intervals of those extreme events also exhibit long-term memory.




FIG. 4: (color online). Detrended fluctuation analysis of the return interval time series for different R_Q . The inset shows the exponential decay of the Hurst index H against R_Q .

We also apply the MFDFA to investigate the multifractal nature of the return intervals. Figure 5(a) shows the power-law dependence of the overall fluctuation $F_q(s)$ on the scale s for $R_Q = 50$. When R_Q increases, the

$F_q(s)$ function becomes more noisy, especially for negative q . The slopes of the straight lines are the linear least-squares estimates of the generalized Hurst indexes $h(q)$, which are drawn in Fig. 5(b). The mass scaling exponent function $\tau(q) = qh(q) - 1$ and the multifractal spectrum $f(\alpha)$ calculated according to the Legendre transform of $\tau(q)$ are illustrated respectively in Fig. 5(c) and Fig. 5(d). The sound nonlinearity in $h(q)$ and $\tau(q)$ is a hallmark of multifractality in the return intervals, whose singularity strength increases with R_Q .




FIG. 5: (color online). Multifractal detrended fluctuation analysis of the return interval time series for different R_Q . (a) MFDFA fluctuation function for different q when $R_Q = 50$. (b) Generalized Hurst indexes $h(q)$. (c) Mass scaling exponents $\tau(q)$. (d) Multifractal singularity spectra $f(\alpha)$.

In summary, we have studied the statistical properties of return intervals of the energy dissipation rate in three-dimensional fully developed turbulence. The interval distribution is found to exhibit scaling behavior across different R_Q and two power-law regimes, except for intervals $r = 1$. We found that the conditional interval distributions also collapse onto a single curve for same r_0 , but deviate for different r_0 , and the mean conditional interval increases as a power law of r_0 , which indicates the presence of short memory in the return intervals. The long-term memory and the multifractal nature in the return intervals are confirmed by the DFA and MFDFA. The Hurst index of the return intervals decays exponentially against R_Q , which allows us to predict the asymptotic Hurst index of return intervals between rare extreme events as $H_\infty = 0.639$. Our results signify discrepancy with the numerical results using MRC and MRW models [12], implying that the energy dissipation process can not be modeled using these models.

This work was partially supported by the National Basic Research Program of China (2004CB217703), the Program for Changjiang Scholars and Innovative Research Team in University (IRT0620), and the Program

for New Century Excellent Talents in University (NCET-07-0288).

* Electronic address: wxzhou@ecust.edu.cn

- [1] A. Bunde, J. Kropp, and H.-J. Schellnhuber, eds., *The Science of Disasters* (Springer, Berlin, 2005).
- [2] K. Y. Kondratyev, C. A. Varotsos, and V. F. Krapivin, *Natural Disasters as Interactive Components of Global Ecodynamics* (Springer, Berlin, 2006).
- [3] M. I. Bogachev, J. F. Eichner, and A. Bunde, *Pure Appl. Geophys.* **165**, 1195 (2008).
- [4] A. Bunde, J. F. Eichner, S. Havlin, and J. W. Kantelhardt, *Physica A* **330**, 1 (2003).
- [5] A. Bunde, J. F. Eichner, J. W. Kantelhardt, and S. Havlin, *Phys. Rev. Lett.* **94**, 048701 (2005).
- [6] E. G. Altmann and H. Kantz, *Phys. Rev. E* **71**, 056106 (2005).
- [7] P. Olla, *Phys. Rev. E* **76**, 011122 (2007).
- [8] M. S. Santhanam and H. Kantz, *Phys. Rev. E* **78**, 051113 (2008).
- [9] J. F. Eichner, J. W. Kantelhardt, A. Bunde, and S. Havlin, *Phys. Rev. E* **75**, 011128 (2007).
- [10] C. Meneveau and K. R. Sreenivasan, *Phys. Rev. Lett.* **59**, 1424 (1987).
- [11] E. Bacry, J. Delour, and J.-F. Muzy, *Phys. Rev. E* **64**, 026103 (2001).
- [12] M. I. Bogachev, J. F. Eichner, and A. Bunde, *Phys. Rev. Lett.* **99**, 240601 (2007).
- [13] M. I. Bogachev, J. F. Eichner, and A. Bunde, *Eur. Phys. J. Spec. Top.* **161**, 181 (2008).
- [14] M. I. Bogachev and A. Bunde, *Phys. Rev. E* **78**, 036114 (2008).
- [15] J.-F. Muzy, J. Delour, and E. Bacry, *Eur. Phys. J. B* **17**, 537 (2000).
- [16] E. Bacry, J. Delour, and J.-F. Muzy, *Physica A* **299**, 84 (2001).
- [17] K. Yamasaki, L. Muchnik, S. Havlin, A. Bunde, and H. E. Stanley, *Proc. Natl. Acad. Sci. USA* **102**, 9424 (2005).
- [18] F.-Z. Wang, K. Yamasaki, S. Havlin, and H. E. Stanley, *Phys. Rev. E* **73**, 026117 (2006).
- [19] F. Wang, P. Weber, K. Yamasaki, S. Havlin, and H. E. Stanley, *Eur. Phys. J. B* **55**, 123 (2007).
- [20] W.-S. Jung, F.-Z. Wang, S. Havlin, T. Kaizoji, H. T. Moon, and H. E. Stanley, *Eur. Phys. J. B* **62**, 113 (2008).
- [21] T. Qiu, L. Guo, and G. Chen, *Physica A* **387**, 6812 (2008).
- [22] F.-Z. Wang, K. Yamasaki, S. Havlin, and H. E. Stanley, *Phys. Rev. E* **77**, 016109 (2008).
- [23] J. W. Lee, K. E. Lee, and P. A. Rikvold, *J. Korean Phys. Soc.* **48**, S123 (2006).
- [24] F. Ren and W.-X. Zhou, *EPL* **84**, 68001 (2008).
- [25] F. Ren, L. Guo, and W.-X. Zhou, *Physica A* **388**, XXX (2009).
- [26] F. Anselmetti, Y. Gagne, E. J. Hopfinger, and R. A. Antonia, *J. Fluid Mech.* **140**, 63 (1984).
- [27] W.-X. Zhou and D. Sornette, *Physica D* **165**, 94 (2002).
- [28] C.-K. Peng, S. V. Buldyrev, S. Havlin, M. Simons, H. E. Stanley, and A. L. Goldberger, *Phys. Rev. E* **49**, 1685 (1994).
- [29] J. W. Kantelhardt, E. Koscielny-Bunde, H. H. A. Rego,

- S. Havlin, and A. Bunde, Physica A **295**, 441 (2001).
[30] J. W. Kantelhardt, S. A. Zschiegner, E. Koscielny-Bunde,
S. Havlin, A. Bunde, and H. E. Stanley, Physica A **316**,
87 (2002).